



Stoller Enterprises, Inc.

VIRO - X

Zn 2.4%, Mn 2.4%, B 1.2%, Cu 1.2%, Mo 0.30% (w.v)

With polyamines and natural plant extracts.

Viro-X is a nutritional product which will increase yield on virus infected plants.

Viruses are protein substances that are vectored into plants by “sucking” insects. The main feeding stimulus of “sucking” insects are amino acids. These compounds are present in newer leaves. The major compounds which allow the virus protein to multiply are other amino acids and nucleotides. This hostile protein (virus) is the cause of plant sickness and possible death.

If viruses encounter “unfamiliar” amino acids (polyamines) they may not be able to replicate. If high levels of divalent metals interfere with enzymes that help the viral protein develop, the virus would not be able to replicate.

Viro-X is designed to do the above. Repeat applications may be necessary on infected plants.

For best results **Viro-X** should be applied with **Stand seedling solution**.

Recommendation:

Apply 9 lt/ha of **Viro-X** when viral infections are first noted.

Apply 9 lt/ha of **Stand** in combination with **Viro-X**.

Use enough water in spray solution to ensure thorough coverage of foliage.

After application, observe any infection in new growth. Symptoms will not disappear from old growth. If symptoms still appear on new growth, repeat application 10 days after first one. Repeat applications 10 days after the second one, if needed.

DO NOT APPLY MORE THAN FOUR APPLICATIONS PER CROP.

